

**SYNTHETIC EXAMPLE**

Alcohol Use Disorders Identification Test

AUDIT - presentation preview

CLIENT

John Doe

ADMINISTERED

Example date

RESPONSE MODE

Self-report

INSTRUMENT

AUDIT

SOURCE

**World Health
Organization**

CLINICAL ROLE

**Alcohol screening
support for clinician
review**

What this report lays out

- Named instrument, version, and provenance
- Item-level response pattern without source wording
- Methodology, references, and clinical boundary
- Responsible clinician review rather than automated conclusions

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored AUDIT result, cannot establish a diagnosis, and must not be placed in a clinical record.

AUDIT does not diagnose alcohol use disorder or determine detox need. Those are clinical decisions.



A report designed for clinical review

Clear structure, visible provenance, and an explicit boundary help a clinician review recorded evidence without mistaking a questionnaire for a conclusion.

REVIEW

Adults complete ten items. Positive findings should trigger fuller assessment and local alcohol-care pathways, including safety review where relevant.

CONTEXT

WHO scoring guidance exists. This page does not independently verify zone labels or production scoring. Use the official manual.

SOURCE

Use WHO materials and local protocol. This catalogue entry does not assign AUDIT from the public site.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.



Item-level layout without source wording

This marketing sample uses invented item numbers and response labels only. It does not reproduce AUDIT item wording, and the pattern below is not a real result.

ITEM	ILLUSTRATIVE SCALE POINT	FICTIONAL RESPONSE
1	3 of 5	Sometimes
2	4 of 5	Often
3	1 of 5	Not at all
4	1 of 5	Not at all
5	1 of 5	Not at all
6	3 of 5	Sometimes
7	3 of 5	Sometimes
8	2 of 5	A little
9	2 of 5	A little
10	5 of 5	Most days
11	4 of 5	Often
12	5 of 5	Most days
13	2 of 5	A little
14	3 of 5	Sometimes

Item numbers, scale points, and responses on this page are invented for layout demonstration. They are not this instrument's item set, scoring structure, or thresholds.



Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The full AUDIT includes consumption items plus questions about control, morning drinking, guilt, blackouts, injury, and concern from others. It is still a screen, not a specialist dependence assessment.

ADMINISTRATION

Adults complete ten items. Positive findings should trigger fuller assessment and local alcohol-care pathways, including safety review where relevant.

CLINICAL BOUNDARY

AUDIT does not diagnose alcohol use disorder or determine detox need. Those are clinical decisions.

GOVERNANCE

Use WHO materials and local protocol. This catalogue entry does not assign AUDIT from the public site.

SELECTED REFERENCES

1. WHO AUDIT manual

World Health Organization. AUDIT: the Alcohol Use Disorders Identification Test.

SOURCE AND VERIFICATION Confirm the official form with the source before any live workflow.