

**SYNTHETIC EXAMPLE**

Drug Abuse Screening Test - 10 item

DAST-10 - presentation preview

CLIENT

John Doe

ADMINISTERED

Example date

RESPONSE MODE

Self-report

INSTRUMENT

DAST-10

SOURCE

**Skinner and subsequent
publishers**

CLINICAL ROLE

**Drug-use screening
support**

What this report lays out

- Named instrument, version, and provenance
- Item-level response pattern without source wording
- Methodology, references, and clinical boundary
- Responsible clinician review rather than automated conclusions

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored DAST-10 result, cannot establish a diagnosis, and must not be placed in a clinical record.

DAST-10 does not diagnose a substance use disorder, identify which drug is involved, or determine treatment.



A report designed for clinical review

Clear structure, visible provenance, and an explicit boundary help a clinician review recorded evidence without mistaking a questionnaire for a conclusion.

REVIEW

Adults complete ten yes/no items. Positive screens should trigger fuller assessment, local policy, and safety review. This page does not collect those answers.

CONTEXT

Published scoring counts endorsed items. This page does not independently verify cutoffs. Confirm the official item text.

SOURCE

Independent verification evidence must be approved before this entry can be assigned.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.



Item-level layout without source wording

This marketing sample uses invented item numbers and response labels only. It does not reproduce DAST-10 item wording, and the pattern below is not a real result.

ITEM	ILLUSTRATIVE SCALE POINT	FICTIONAL RESPONSE
1	2 of 5	A little
2	5 of 5	Most days
3	3 of 5	Sometimes
4	3 of 5	Sometimes
5	4 of 5	Often
6	1 of 5	Not at all
7	1 of 5	Not at all
8	5 of 5	Most days
9	5 of 5	Most days
10	5 of 5	Most days
11	3 of 5	Sometimes
12	3 of 5	Sometimes
13	2 of 5	A little
14	1 of 5	Not at all

Item numbers, scale points, and responses on this page are invented for layout demonstration. They are not this instrument's item set, scoring structure, or thresholds.



Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Drug Abuse Screening Test-10 asks about drug-related problems excluding alcohol in many standard instructions. It is a brief screen, not a toxicology result and not a full substance-use interview.

ADMINISTRATION

Adults complete ten yes/no items. Positive screens should trigger fuller assessment, local policy, and safety review. This page does not collect those answers.

CLINICAL BOUNDARY

DAST-10 does not diagnose a substance use disorder, identify which drug is involved, or determine treatment.

GOVERNANCE

Independent verification evidence must be approved before this entry can be assigned.

SELECTED REFERENCES

1. DAST development

Skinner HA. The drug abuse screening test. *Addict Behav.* 1982;7(4):363-371.

SOURCE AND VERIFICATION Confirm the official form with the source before any live workflow.