



SYNTHETIC EXAMPLE

Demoralization Scale-II

DS-II - presentation preview

CLIENT

John Doe

ADMINISTERED

Example date

RESPONSE MODE

Self-report

INSTRUMENT

DS-II

SOURCE

Robinson, Kissane and
colleagues

CLINICAL ROLE

Demoralization
screening support for
clinician review

What this report lays out

- Named instrument, version, and provenance
- Item-level response pattern without source wording
- Methodology, references, and clinical boundary
- Responsible clinician review rather than automated conclusions

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored DS-II result, cannot establish a diagnosis, and must not be placed in a clinical record.

DS-II is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.



A report designed for clinical review

Clear structure, visible provenance, and an explicit boundary help a clinician review recorded evidence without mistaking a questionnaire for a conclusion.

REVIEW

DS-II is completed as a focused self-report and stored for clinician review. This public page does not collect answers or return a score.

CONTEXT

Published scoring conventions may exist in official materials. This public page does not independently verify scoring, cutoffs, or a production scoring rule. Use the official form and scoring instructions for the version you intend to use.

SOURCE

Catalogue information on the measure's purpose, source and administration requirements. Check the official administration guidance and suitability for your clinical purpose.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.



Item-level layout without source wording

This marketing sample uses invented item numbers and response labels only. It does not reproduce DS-II item wording, and the pattern below is not a real result.

ITEM	ILLUSTRATIVE SCALE POINT	FICTIONAL RESPONSE
1	3 of 5	Sometimes
2	1 of 5	Not at all
3	3 of 5	Sometimes
4	2 of 5	A little
5	1 of 5	Not at all
6	2 of 5	A little
7	2 of 5	A little
8	1 of 5	Not at all
9	2 of 5	A little
10	4 of 5	Often
11	1 of 5	Not at all
12	3 of 5	Sometimes
13	2 of 5	A little
14	2 of 5	A little

Item numbers, scale points, and responses on this page are invented for layout demonstration. They are not this instrument's item set, scoring structure, or thresholds.



Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Demoralization Scale-II asks about meaning, purpose, and coping over the past two weeks. Screening evidence around a total of 20 is a prompt for clinician review. It is not a diagnosis and not an end-of-life plan.

ADMINISTRATION

DS-II is completed as a focused self-report and stored for clinician review. This public page does not collect answers or return a score.

CLINICAL BOUNDARY

DS-II is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

GOVERNANCE

Catalogue information on the measure's purpose, source and administration requirements. Check the official administration guidance and suitability for your clinical purpose.

SELECTED REFERENCES

1. DS-II source literature

Robinson S, Kissane DW et al. Demoralization Scale-II source literature used by this catalogue entry.

SOURCE AND VERIFICATION Confirm the official form with the source before any live workflow.