



SYNTHETIC EXAMPLE

Edinburgh Postnatal Depression Scale

EPDS - presentation preview

CLIENT

John Doe

ADMINISTERED

Example date

RESPONSE MODE

Self-report

INSTRUMENT

EPDS

SOURCE

Cox, Holden and
Sagovsky

CLINICAL ROLE

Perinatal screening
support for clinician
review

What this report lays out

- Named instrument, version, and provenance
- Item-level response pattern without source wording
- Methodology, references, and clinical boundary
- Responsible clinician review rather than automated conclusions

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored EPDS result, cannot establish a diagnosis, and must not be placed in a clinical record.

EPDS does not diagnose postnatal depression, anxiety, or psychosis. A low score does not rule out illness. Any concerning item requires clinician-led review.



A report designed for clinical review

Clear structure, visible provenance, and an explicit boundary help a clinician review recorded evidence without mistaking a questionnaire for a conclusion.

REVIEW

The person completes ten items about the past seven days. Results should reach a clinician who understands perinatal context, social support, and local safeguarding. Lirena does not provide urgent monitoring.

CONTEXT

Published scoring and suggested thresholds vary by setting and language. This page does not independently verify a cutoff. Use the official version, timing, and local protocol.

SOURCE

Confirm the translation you use, the timing relative to birth or pregnancy, and your review pathway. This catalogue entry does not assign EPDS from the public site.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.



Item-level layout without source wording

This marketing sample uses invented item numbers and response labels only. It does not reproduce EPDS item wording, and the pattern below is not a real result.

ITEM	ILLUSTRATIVE SCALE POINT	FICTIONAL RESPONSE
1	1 of 5	Not at all
2	2 of 5	A little
3	4 of 5	Often
4	4 of 5	Often
5	3 of 5	Sometimes
6	1 of 5	Not at all
7	3 of 5	Sometimes
8	2 of 5	A little
9	4 of 5	Often
10	3 of 5	Sometimes
11	3 of 5	Sometimes
12	4 of 5	Often
13	2 of 5	A little
14	4 of 5	Often

Item numbers, scale points, and responses on this page are invented for layout demonstration. They are not this instrument's item set, scoring structure, or thresholds.



Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The EPDS asks about how a pregnant or postnatal person has felt in the past week. It is widely used as a perinatal screening aid. Item-level review, including any self-harm-related response, remains a clinician responsibility.

ADMINISTRATION

The person completes ten items about the past seven days. Results should reach a clinician who understands perinatal context, social support, and local safeguarding. Lirena does not provide urgent monitoring.

CLINICAL BOUNDARY

EPDS does not diagnose postnatal depression, anxiety, or psychosis. A low score does not rule out illness. Any concerning item requires clinician-led review.

GOVERNANCE

Confirm the translation you use, the timing relative to birth or pregnancy, and your review pathway. This catalogue entry does not assign EPDS from the public site.

SELECTED REFERENCES

1. Original EPDS paper

Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Br J Psychiatry. 1987;150:782-786.

SOURCE AND VERIFICATION Confirm the official form with the source before any live workflow.