

**SYNTHETIC EXAMPLE**

Montgomery-Asberg Depression Rating Scale

MADRS - presentation preview

CLIENT

John Doe

ADMINISTERED

Example date

RESPONSE MODE

Clinician-rated

INSTRUMENT

MADRS

SOURCE

Montgomery and Asberg

CLINICAL ROLE

**Clinician-rated
severity support**

What this report lays out

- Named instrument, version, and provenance
- Item-level response pattern without source wording
- Methodology, references, and clinical boundary
- Responsible clinician review rather than automated conclusions

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored MADRS result, cannot establish a diagnosis, and must not be placed in a clinical record.

A MADRS total is a clinician's rated severity estimate, not a diagnosis and not a treatment recommendation. Interview quality, rater training, and clinical context all matter.



A report designed for clinical review

Clear structure, visible provenance, and an explicit boundary help a clinician review recorded evidence without mistaking a questionnaire for a conclusion.

REVIEW

A clinician interviews the person and rates each item against published anchors. This is not a client-only form. Digital workflows must preserve the rater's role and the official rating instructions.

CONTEXT

Published manuals describe item anchors and total-score ranges. This page does not independently verify a scoring rule or cutoff. Use the official manual for the version your service intends to use.

SOURCE

Check the exact form and whether your raters are trained for the version you want. This catalogue entry does not assign MADRS from the public site.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.



Item-level layout without source wording

This marketing sample uses invented item numbers and response labels only. It does not reproduce MADRS item wording, and the pattern below is not a real result.

ITEM	ILLUSTRATIVE SCALE POINT	FICTIONAL RESPONSE
1	1 of 5	Not at all
2	4 of 5	Often
3	3 of 5	Sometimes
4	5 of 5	Most days
5	2 of 5	A little
6	5 of 5	Most days
7	2 of 5	A little
8	3 of 5	Sometimes
9	1 of 5	Not at all
10	1 of 5	Not at all
11	5 of 5	Most days
12	1 of 5	Not at all
13	4 of 5	Often
14	2 of 5	A little

Item numbers, scale points, and responses on this page are invented for layout demonstration. They are not this instrument's item set, scoring structure, or thresholds.



Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

MADRS was designed to be sensitive to change in depression treatment studies. A trained clinician rates defined items after interview. It is not a patient-completed questionnaire and should not be presented as one.

ADMINISTRATION

A clinician interviews the person and rates each item against published anchors. This is not a client-only form. Digital workflows must preserve the rater's role and the official rating instructions.

CLINICAL BOUNDARY

A MADRS total is a clinician's rated severity estimate, not a diagnosis and not a treatment recommendation. Interview quality, rater training, and clinical context all matter.

GOVERNANCE

Check the exact form and whether your raters are trained for the version you want. This catalogue entry does not assign MADRS from the public site.

SELECTED REFERENCES

1. Original MADRS paper

Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. Br J Psychiatry. 1979;134:382-389.

SOURCE AND VERIFICATION Confirm the official form with the source before any live workflow.