

**SYNTHETIC EXAMPLE**

Positive and Negative Syndrome Scale

PANSS - presentation preview

CLIENT

John Doe

ADMINISTERED

Example date

RESPONSE MODE

Clinician-rated

INSTRUMENT

PANSS

SOURCE

**Kay, Fiszbein and Opler
/ current publisher**

CLINICAL ROLE

**Specialist clinician
rating**

What this report lays out

- Named instrument, version, and provenance
- Item-level response pattern without source wording
- Methodology, references, and clinical boundary
- Responsible clinician review rather than automated conclusions

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored PANSS result, cannot establish a diagnosis, and must not be placed in a clinical record.

PANSS does not diagnose schizophrenia. It is a specialist severity rating inside clinician-led care.



A report designed for clinical review

Clear structure, visible provenance, and an explicit boundary help a clinician review recorded evidence without mistaking a questionnaire for a conclusion.

REVIEW

A trained rater uses official items and anchors. This cannot be converted into a public questionnaire.

CONTEXT

Official manuals define subscale scoring. This page does not independently verify algorithms or reproduce items.

SOURCE

Independent verification evidence must be approved before this entry can be assigned.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.



Item-level layout without source wording

This marketing sample uses invented item numbers and response labels only. It does not reproduce PANSS item wording, and the pattern below is not a real result.

ITEM	ILLUSTRATIVE SCALE POINT	FICTIONAL RESPONSE
1	5 of 5	Most days
2	5 of 5	Most days
3	5 of 5	Most days
4	2 of 5	A little
5	1 of 5	Not at all
6	2 of 5	A little
7	5 of 5	Most days
8	4 of 5	Often
9	3 of 5	Sometimes
10	5 of 5	Most days
11	3 of 5	Sometimes
12	5 of 5	Most days
13	4 of 5	Often
14	3 of 5	Sometimes

Item numbers, scale points, and responses on this page are invented for layout demonstration. They are not this instrument's item set, scoring structure, or thresholds.



Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Positive and Negative Syndrome Scale is used in specialist psychosis services and trials to rate positive, negative, and general psychopathology items after interview and observation.

ADMINISTRATION

A trained rater uses official items and anchors. This cannot be converted into a public questionnaire.

CLINICAL BOUNDARY

PANSS does not diagnose schizophrenia. It is a specialist severity rating inside clinician-led care.

GOVERNANCE

Independent verification evidence must be approved before this entry can be assigned.

SELECTED REFERENCES

1. Original PANSS paper

Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS). Schizophr Bull. 1987;13(2):261-276.

SOURCE AND VERIFICATION Confirm the official form with the source before any live workflow.