

**SYNTHETIC EXAMPLE**

Young Mania Rating Scale

YMRS - presentation preview

CLIENT**John Doe****ADMINISTERED****Example date****RESPONSE MODE****Clinician-rated****INSTRUMENT****YMRS****SOURCE****Young and colleagues****CLINICAL ROLE****Clinician-rated mania
severity support**

What this report lays out

- Named instrument, version, and provenance
- Item-level response pattern without source wording
- Methodology, references, and clinical boundary
- Responsible clinician review rather than automated conclusions

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored YMRS result, cannot establish a diagnosis, and must not be placed in a clinical record.

A YMRS rating cannot diagnose mania, specify bipolar subtype, or determine treatment. It is one severity estimate inside clinician-led care.



A report designed for clinical review

Clear structure, visible provenance, and an explicit boundary help a clinician review recorded evidence without mistaking a questionnaire for a conclusion.

REVIEW

A clinician rates items after interview and observation. It should not be converted into an unsupervised questionnaire. Digital capture must keep the rater in the loop.

CONTEXT

Published item anchors and totals exist in the source paper and later manuals. This page does not independently verify scoring. Use the official rating instructions.

SOURCE

Confirm rater training and the exact form. This catalogue entry does not assign YMRS from the public site.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.



Item-level layout without source wording

This marketing sample uses invented item numbers and response labels only. It does not reproduce YMRS item wording, and the pattern below is not a real result.

ITEM	ILLUSTRATIVE SCALE POINT	FICTIONAL RESPONSE
1	4 of 5	Often
2	1 of 5	Not at all
3	1 of 5	Not at all
4	2 of 5	A little
5	5 of 5	Most days
6	4 of 5	Often
7	5 of 5	Most days
8	1 of 5	Not at all
9	4 of 5	Often
10	3 of 5	Sometimes
11	3 of 5	Sometimes
12	2 of 5	A little
13	3 of 5	Sometimes
14	3 of 5	Sometimes

Item numbers, scale points, and responses on this page are invented for layout demonstration. They are not this instrument's item set, scoring structure, or thresholds.



Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

YMRS is used to rate manic symptom severity based on clinician observation and interview. It is common in research and some specialist mood services. It does not diagnose bipolar disorder by itself.

ADMINISTRATION

A clinician rates items after interview and observation. It should not be converted into an unsupervised questionnaire. Digital capture must keep the rater in the loop.

CLINICAL BOUNDARY

A YMRS rating cannot diagnose mania, specify bipolar subtype, or determine treatment. It is one severity estimate inside clinician-led care.

GOVERNANCE

Confirm rater training and the exact form. This catalogue entry does not assign YMRS from the public site.

SELECTED REFERENCES

1. Original YMRS paper

Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: reliability, validity and sensitivity. Br J Psychiatry. 1978;133:429-435.

SOURCE AND VERIFICATION Confirm the official form with the source before any live workflow.